



# Image alignment



# A look into the past



<http://blog.flickr.net/en/2010/01/27/a-look-into-the-past/>

# A look into the past

- Leningrad during the blockade




<http://komen-dant.livejournal.com/345684.html>

# Bing streetside images




<http://www.bing.com/community/blogs/maps/archive/2010/01/12/new-bing-maps-application-streetside-photos.aspx>

# Image alignment: Applications




Panorama stitching




Recognition of object instances

# Image alignment: Challenges




Small degree of overlap


Intensity changes




Occlusion,  
clutter



# Image alignment




- Two families of approaches:
  - **Direct (pixel-based) alignment**
    - Search for alignment where most pixels agree
  - **Feature-based alignment**
    - Search for alignment where *extracted features* agree
    - Can be verified using pixel-based alignment



# Alignment as fitting

- Previous lectures: fitting a model to features in one image




Find model  $M$  that minimizes

$$\sum_i \text{residual}(x_i, M)$$

# Alignment as fitting



Previous lectures: fitting a model to features in one image



Find model  $M$  that minimizes

$$\sum_i \text{residual}(x_i, M)$$

- Alignment: fitting a model to a transformation between pairs of features (*matches*) in two images




Find transformation  $T$  that minimizes


$$\sum_i \text{residual}(T(x_i), x'_i)$$

# 2D transformation models


- Similarity  
(translation, scale, rotation)



- Affine




- Projective  
(homography)




# Let's start with affine transformations

- Simple fitting procedure (linear least squares)
- Approximates viewpoint changes for roughly planar objects and roughly orthographic cameras
- Can be used to initialize fitting for more complex models




# Fitting an affine transformation

- Assume we know the correspondences, how do we get the transformation?



$$\begin{bmatrix} x'_i \\ y'_i \end{bmatrix} = \begin{bmatrix} m_1 & m_2 \\ m_3 & m_4 \end{bmatrix} \begin{bmatrix} x_i \\ y_i \end{bmatrix} + \begin{bmatrix} t_1 \\ t_2 \end{bmatrix}$$

$$\begin{bmatrix} \dots \\ x_i & y_i & 0 & 0 & 1 & 0 \\ 0 & 0 & x_i & y_i & 0 & 1 \\ \dots \end{bmatrix} \begin{bmatrix} m_1 \\ m_2 \\ m_3 \\ m_4 \\ t_1 \\ t_2 \end{bmatrix} = \begin{bmatrix} \dots \\ x'_i \\ y'_i \\ \dots \end{bmatrix}$$




## Fitting an affine transformation

---

$$\begin{bmatrix} \dots & & & & & & \\ x_i & y_i & 0 & 0 & 1 & 0 & \\ 0 & 0 & x_i & y_i & 0 & 1 & \\ \dots & & & & & & \end{bmatrix} \begin{bmatrix} m_1 \\ m_2 \\ m_3 \\ m_4 \\ t_1 \\ t_2 \end{bmatrix} = \begin{bmatrix} \dots \\ x'_i \\ y'_i \\ \dots \end{bmatrix}$$


- Linear system with six unknowns
- Each match gives us two linearly independent equations: need at least three to solve for the transformation parameters



# Fitting a plane projective transformation

---

- **Homography:** plane projective transformation (transformation taking a quad to another arbitrary quad)




# Homography



- The transformation between two views of a planar surface




- The transformation between images from two cameras that share the same center



# Application: Panorama stitching



Source: Hartley & Zisserman



# Fitting a homography

---


- Recall: homogeneous coordinates

$$(x, y) \Rightarrow \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

Converting *to* homogeneous  
image coordinates

$$\begin{bmatrix} x \\ y \\ w \end{bmatrix} \Rightarrow (x/w, y/w)$$

Converting *from* homogeneous  
image coordinates



# Fitting a homography

- Recall: homogeneous coordinates

$$(x, y) \Rightarrow \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} \qquad \begin{bmatrix} x \\ y \\ w \end{bmatrix} \Rightarrow (x/w, y/w)$$

Converting *to* homogeneous  
image coordinates

Converting *from* homogeneous  
image coordinates

- Equation for homography:

$$\lambda \begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} h_{11} & h_{12} & h_{13} \\ h_{21} & h_{22} & h_{23} \\ h_{31} & h_{32} & h_{33} \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

# Fitting a homography

- Equation for homography:

$$\lambda \begin{bmatrix} x'_i \\ y'_i \\ 1 \end{bmatrix} = \begin{bmatrix} h_{11} & h_{12} & h_{13} \\ h_{21} & h_{22} & h_{23} \\ h_{31} & h_{32} & h_{33} \end{bmatrix} \begin{bmatrix} x_i \\ y_i \\ 1 \end{bmatrix} \quad \lambda \mathbf{x}'_i = \mathbf{H} \mathbf{x}_i$$

$$\mathbf{x}'_i \times \mathbf{H} \mathbf{x}_i = 0$$

$$\begin{bmatrix} x'_i \\ y'_i \\ 1 \end{bmatrix} \times \begin{bmatrix} \mathbf{h}_1^T \mathbf{x}_i \\ \mathbf{h}_2^T \mathbf{x}_i \\ \mathbf{h}_3^T \mathbf{x}_i \end{bmatrix} = \begin{bmatrix} y'_i \mathbf{h}_3^T \mathbf{x}_i - \mathbf{h}_2^T \mathbf{x}_i \\ \mathbf{h}_1^T \mathbf{x}_i - x'_i \mathbf{h}_3^T \mathbf{x}_i \\ x'_i \mathbf{h}_2^T \mathbf{x}_i - y'_i \mathbf{h}_1^T \mathbf{x}_i \end{bmatrix}$$

$$\begin{bmatrix} 0^T & -\mathbf{x}_i^T & y'_i \mathbf{x}_i^T \\ \mathbf{x}_i^T & 0^T & -x'_i \mathbf{x}_i^T \\ -y'_i \mathbf{x}_i^T & x'_i \mathbf{x}_i^T & 0^T \end{bmatrix} \begin{pmatrix} \mathbf{h}_1 \\ \mathbf{h}_2 \\ \mathbf{h}_3 \end{pmatrix} = 0$$

3 equations,  
only 2 linearly  
independent

## Direct linear transform

$$\begin{bmatrix} 0^T & \mathbf{x}_1^T & -y'_1 \mathbf{x}_1^T \\ \mathbf{x}_1^T & 0^T & -x'_1 \mathbf{x}_1^T \\ \dots & \dots & \dots \\ 0^T & \mathbf{x}_n^T & -y'_n \mathbf{x}_n^T \\ \mathbf{x}_n^T & 0^T & -x'_n \mathbf{x}_n^T \end{bmatrix} \begin{pmatrix} \mathbf{h}_1 \\ \mathbf{h}_2 \\ \mathbf{h}_3 \end{pmatrix} = 0 \quad \mathbf{A} \mathbf{h} = 0$$

- H has 8 degrees of freedom (9 parameters, but scale is arbitrary)
- One match gives us two linearly independent equations
- Four matches needed for a minimal solution (null space of 8x9 matrix)
- More than four: homogeneous least squares


# Example



1st view



2nd view



2nd view warped  
by the planar homography  
between two views

# Rotation Only - Calibrated Case

- Calibrated Two views related by rotation only

$$\lambda_2 \mathbf{x}_2 = R \lambda_1 \mathbf{x}_1 \quad \widehat{\mathbf{x}}_2 R \mathbf{x}_1 = 0$$

- Mapping to a reference view – rotation can be estimated



- Mapping to a cylindrical surface

